

A note on the classification of naturally graded Lie algebras with linear characteristic sequence

José María Ancochea Bermúdez*

Rutwig Campoamor†

Departamento de Geometría y Topología
Fac. CC. Matemáticas Univ. Complutense
28040 Madrid (Spain)

Abstract

For sufficiently high dimensions, the naturally graded nonsplit nilpotent Lie algebras with linear characteristic sequence are classified.

Introduction

The determination by Vergne of the naturally graded filiform Lie algebras [6] in relation with the study of irreducible components of the variety of nilpotent Lie algebra laws pointed out the interest of graded structures for this class of algebras. The introduction of the characteristic sequence of a nilpotent Lie algebra [1] made it feasible to formulate the problem for nilpotent Lie algebras of lower nilindexes.

Computational methods have been applied in [5] to determine the algebras of characteristic sequence $(n - 3, 1, 1, 1)$ for $n \geq 8$. With few exceptions (due to the weakness of the conditions imposed by the graduation in low dimensions) these algebras belong to a couple of families. However, the weight of this procedure relies heavily on the computations, which restricts the generality of the results obtained. Following the philosophy of [6], i.e., using both extension theory and the cohomology spaces with trivial coefficients, the problem can be satisfactorily solved in arbitrary dimension and nilindexes, up to a reduced number of pathological cases appearing in low dimensions. In this sense, all the naturally graded Lie algebras of characteristic sequence $(2m - 1, q, 1)$ with $m \geq 3$ and $1 \leq q \leq 2m - 3$ were classified in [2].

Finally. as a global classification of nilpotent Lie algebras is a hopeless problem [4], it seems more reasonable to analyze the graded structures and then use deformation theory to obtain concrete isomorphism classes satisfying certain

*corresponding author: e-mail: Jose_Ancochea@mat.ucm.es

†e-mail: rutwig@nfssrv.mat.ucm.es

properties. This has for example been done to obtain nilradicals of complete and rigid Lie algebras [3].

1 Preliminaries and notations

Whenever we speak about Lie algebras in this work, we refer to finite dimensional complex nonsplit Lie algebras.

Let $\mathfrak{g}$ be an n -dimensional nilpotent Lie algebra. We denote by $C^i(\mathfrak{g})$ the ideals of the central descending sequence of $\mathfrak{g}$. Then the $\mathbb{Z}$ -graded Lie algebra associated to this filtration is defined by

$$\mathfrak{gr}(\mathfrak{g}) = \sum_{i \in \mathbb{Z}} \frac{C^i \mathfrak{g}}{C^{i+1} \mathfrak{g}}$$

Definition. A nilpotent Lie algebra $\mathfrak{g}$ is called naturally graded if it is isomorphic to $\mathfrak{gr}(\mathfrak{g})$.

In [1] the following invariant is presented :

For any nonzero vector $X \in \mathfrak{g}_n - C^1 \mathfrak{g}_n$ let $c(X)$ be the ordered sequence of a similitude invariant for the nilpotent operator $ad_\mu(X)$; i.e., the ordered sequence of dimensions of Jordan blocks of this operator. The set of these sequences is ordered lexicographically.

Definition. The characteristic sequence of $\mathfrak{g}_n$ is an isomorphism invariant $c(\mathfrak{g}_n)$ defined by

$$c(\mathfrak{g}_n) = \max_{X \in \mathfrak{g}_n - C^1 \mathfrak{g}_n} \{c(X)\}$$

A nonzero vector $X \in \mathfrak{g}_n - C^1 \mathfrak{g}_n$ for which $c(X) = c(\mathfrak{g}_n)$ is called characteristic vector.

A characteristic sequence is called linear if it is of the form $(q, 1, \dots, 1)$ for $q \geq 2$.

A Lie algebra $\mathfrak{g}$ is called filiform if $c(\mathfrak{g}) = (n - 1, 1)$, where n is the dimension. In [6], Vergne proves that every naturally graded filiform Lie algebra is isomorphic to one of the algebras L_n and Q_n defined by

1. L_n ($n \geq 3$) :

$$[X_1, X_i] = X_{i+1}, \quad 1 \leq i \leq n$$

over the basis $\{X_1, \dots, X_{n+1}\}$.

2. Q_{2m-1} ($m \geq 3$) :

$$[X_1, X_i] = X_{i+1}, \quad 1 \leq i \leq 2m - 1$$

$$[X_j, X_{2m+1-j}] = (-1)^j X_{2m}, \quad 1 \leq j \leq m$$

over the basis $\{X_1, \dots, X_{2m}\}$.

It will be convenient to use the so called contragradient representation of a Lie algebra $\mathfrak{g}$. Let $n = \dim(\mathfrak{g})$ and $\{X_1, \dots, X_n\}$ be a basis. If $C_{i,j}^k$ are the structure constants of the algebra law μ , we can define, over the dual basis $\{\omega_1, \dots, \omega_n\}$, the differential

$$d_\mu \omega_i(X_j, X_k) = -C_{jk}^i$$

Then the Lie algebra is rewritten as

$$d\omega_i = -C_{jk}^i \omega_j \wedge \omega_k \quad 1 \leq i, j, k \leq n,$$

The Jacobi condition is equivalent to $d^2\omega_i = 0$ for all i . Thus the problem of determining the structure constants is translated to the calculation of a system of closed differential forms. This approach is more useful in practice, as it allows to recognize both extensions and central quotients faster. This also explains why for the first dimensions admitting a certain graduation type and a linear characteristic sequence, the conditions on the structure constants are too weak.

The two central results to classify the naturally graded Lie algebras in dimension n and linear characteristic sequence $(n-t, 1, \dots, 1)$ for $2 \leq t \leq \lfloor \frac{n}{2} \rfloor$ are

Proposition 1. *Let $\mathfrak{g}$ be a split naturally graded Lie algebra of nilindex p and linear characteristic sequence. Then $\mathfrak{g}$ does not admit a nonsplit naturally graded central extension of degree one preserving both the nilindex and the linearity of $c(\mathfrak{g})$.*

Proposition 2. *Let $\mathfrak{g}$ an n -dimensional naturally graded Lie algebra with nilindex p and linear characteristic sequence. Then there exists an $(n-1)$ -dimensional naturally graded Lie algebra $\mathfrak{g}'$ with the same nilindex and linear characteristic sequence such that the following sequence is exact :*

$$0 \rightarrow \mathbb{C} \rightarrow \mathfrak{g} \rightarrow \mathfrak{g}' \rightarrow 0$$

The proof of the first assertion is trivial, as any nonsplit graded extension preserving the nilindex and the linearity of the characteristic sequence must necessarily involcrate the characteristic vectors, which implies the generation of a Jordan block of dimension two for the adjoint operator of such a vector. As to the second proposition, for sufficient large dimensions it is not difficult to see that the conditions on the coefficients of the contragradient representation do not depend any more on the dimension of the algebra, but only on the graduation type [2].

Starting from these auxiliary results and proceeding like in [2] we obtain :

Theorem 1. *For sufficiently high dimension, any naturally graded Lie algebra $\mathfrak{g}$ of characteristic sequence $(\dim \mathfrak{g} - k, 1, \dots, 1)$ with $2 \leq k \leq \lfloor \frac{\dim \mathfrak{g}}{2} \rfloor$ is isomorphic to one of the following models :*

$$1. L_n(t_1, \dots, t_p) ; 1 \leq t_1 < \dots < t_p \leq \left[\frac{n-1}{2} \right], n \geq 7$$

$$\begin{aligned} d\omega_1 &= d\omega_2 = 0 \\ d\omega_j &= \omega_1 \wedge \omega_{j-1}, \quad 3 \leq j \leq n \\ d\omega_{n+i} &= \sum_{j=2}^{t_i+1} (-1)^j \omega_j \wedge \omega_{2t_i+3-j}, \quad 1 \leq i \leq p \end{aligned}$$

$$2. Q_{2m-1}(t_1, \dots, t_p) ; 1 \leq t_1 < \dots < t_p \leq \left[\frac{2m-1}{2} \right], m \geq 3$$

$$\begin{aligned} d\omega_1 &= d\omega_2 = 0 \\ d\omega_j &= \omega_1 \wedge \omega_{j-1}, \quad 3 \leq j \leq 2m-1 \\ d\omega_{2m} &= \omega_1 \wedge \omega_{2m-1} + \sum_{j=2}^{\left[\frac{2m+1}{2} \right]} (-1)^j \omega_j \wedge \omega_{2m+1-j} \\ d\omega_{2m+1+j} &= \sum_{i=2}^{t_j} (-1)^j \omega_i \wedge \omega_{2t_j+3-i}, \quad 1 \leq j \leq p \end{aligned}$$

$$3. D_{2m}(t_1, \dots, t_p) ; 1 \leq t_1 < \dots < t_p \leq m-3, m \geq 4$$

$$\begin{aligned} d\omega_1 &= d\omega_2 = 0 \\ d\omega_j &= \omega_1 \wedge \omega_{j-1}, \quad 3 \leq j \leq 2m-3 \\ d\omega_{2m-2} &= \omega_1 \wedge \omega_{2m-3} + \sum_{j=2}^{m-1} (-1)^j \omega_j \wedge \omega_{2m-3-j} \\ d\omega_{2m-1} &= \omega_1 \wedge \omega_{2m-2} + \sum_{j=2}^{m-1} (-1)^j (m-j) \omega_j \wedge \omega_{2m-2-j} \\ &+ (2-m) \omega_2 \wedge \omega_{2m} \\ d\omega_{2m} &= \sum_{j=2}^{m-1} (-1)^j \omega_j \wedge \omega_{2m-3-j} \\ d\omega_{2m+i} &= \sum_{j=2}^{t_i-1} (-1)^j \omega_j \wedge \omega_{2t_i+3-j}, \quad 1 \leq i \leq p \end{aligned}$$

4. $E_{2m+1}(t_1, \dots, t_p)$; $1 \leq t_1 < \dots < t_p \leq m-3$, $m \geq 4$

$$\begin{aligned}
d\omega_1 &= d\omega_2 = 0 \\
d\omega_j &= \omega_1 \wedge \omega_{j-1}, \quad 3 \leq j \leq 2m-3 \\
d\omega_{2m-2} &= \omega_1 \wedge \omega_{2m-3} + \sum_{j=2}^{m-1} (-1)^j \omega_j \wedge \omega_{2m-1-j} \\
d\omega_{2m-1} &= \omega_1 \wedge \omega_{2m-2} + \sum_{j=2}^{m-1} (-1)^j (m-j) \omega_j \wedge \omega_{2m-j} \\
&+ (m-2) \omega_2 \wedge \omega_{2m+1} \\
d\omega_{2m} &= \omega_1 \wedge \omega_{2m-1} + \sum_{j=3}^m \frac{(-1)^j (j-2)(2m-1-j)}{2} \omega_j \wedge \omega_{2m+1-j} \\
&+ (m-2) \omega_3 \wedge \omega_{2m+1} \\
d\omega_{2m+1} &= \sum_{j=2}^{\left[\frac{2m+1}{2}\right]-1} (-1)^j \omega_j \wedge \omega_{2m-1-j} \\
d\omega_{2m+1+i} &= \sum_{j=2}^{t_i+1} (-1)^j \omega_j \wedge \omega_{2t_i+3-j}, \quad 1 \leq i \leq p
\end{aligned}$$

The result can be resumed in the following table. The fourth column indicates the restrictions on n .

$\mathfrak{g}$	$\dim \mathfrak{g}$	$ch(\mathfrak{g})$	
$L_n(t_1, \dots, t_p)$	$n+p$	$(n-1, 1, \dots, {}^{(p+1)}, \dots, 1)$	
$Q_n(t_1, \dots, t_p)$	$2m+p$	$(2m-1, 1, \dots, {}^{(p+1)}, \dots, 1)$	$n = 2m-1$
$D_n(t_1, \dots, t_p)$	$2m+p$	$(2m-2, 1, \dots, {}^{(p+2)}, \dots, 1)$	$n = 2m$
$E_n(t_1, \dots, t_p)$	$2m+1+p$	$(2m-1, 1, \dots, {}^{(p+2)}, \dots, 1)$	$n = 2m+1$

Corollary 1. *For any dimension $n \geq 7$ the number of naturally graded Lie algebras with linear characteristic sequence is finite.*

References

- [1] J. M. Ancochea, M. Goze. *Sur la classification des algèbres de Lie nilpotentes de dimension 7*, C.R.A.S. 302 (1986), 611-613.
- [2] J. M. Ancochea, R. Campoamor. *On certain families of naturally graded Lie algebras*, J. Pure Appl. Algebra, to appear.
- [3] J. M. Ancochea, R. Campoamor. *On nonfiliform characteristically nilpotent and complete Lie algebras*, Alg. Colloq., to appear.
- [4] L. Y. Galitski, D.A. Timashev. *On classifications of metabelian Lie algebras*, J. of Lie theory 9 (1999), 125-156.

- [5] E. Pastor Sahagún. *Graduaciones naturales de álgebras de Lie 3-filiformes* , Ph.D. Thesis, 2001.
- [6] M. Vergne. *Cohomologie des algèbres de Lie nilpotentes. Applications a l'étude de la variété des algèbres de Lie nilpotentes*, Bull. Soc. Math. France 98 (1970), 81-116.